



LIFE·PAC®

Science



Alpha Omega Publications®

SCIENCE 1209

ATOMIC AND NUCLEAR PHYSICS

CONTENTS

I. QUANTUM THEORY	2
ELECTROMAGNETIC RADIATION.....	2
MATTER WAVES	8
ATOMIC MODELS	11
ATOMIC SPECTRA.....	12
BOHR MODEL.....	14
II. NUCLEAR THEORY	22
BUILDING BLOCKS OF THE NUCLEUS	22
PROPERTIES OF THE NUCLEUS	23
III. NUCLEAR REACTIONS	30
NUCLEAR FISSION	30
FUSION.....	32
NUCLEAR REACTION APPLICATIONS	33
GLOSSARY	39
FORMULAS	41

Author:

Robert Hughson, M.S.

Editor:

Alan Christopherson, M.S.

Illustrations:

Jim Lahowetz

Alpha Omega Graphics



Alpha Omega Publications®

804 N. 2nd Ave. E., Rock Rapids, IA 51246-1759

© MM by Alpha Omega Publications, Inc. All rights reserved.

LIFEPAAC is a registered trademark of Alpha Omega Publications, Inc.

All trademarks and/or service marks referenced in this material are the property of their respective owners. Alpha Omega Publications, Inc. makes no claim of ownership to any trademarks and/or service marks other than their own and their affiliates', and makes no claim of affiliation to any companies whose trademarks may be listed in this material, other than their own.

ATOMIC AND NUCLEAR PHYSICS

You are about to begin the study of modern physics, the physics of very small objects. You will learn more about how atoms and their subparts interact. The laws of physics that you have already learned will be used, but you will notice that at times some new concepts will be considered.

OBJECTIVES

Read these objectives. The objectives tell you what you will be able to do when you have successfully completed this LIFEPAAC®.

When you have finished this LIFEPAAC, you should be able to:

1. Explain when wave theory is applicable and when quantum theory is applicable.
2. Calculate the energy of photoelectrons and X rays.
3. Explain de Broglie waves.
4. State and apply the uncertainty principle.
5. Define emission spectra and absorption spectra.
6. Define line spectra and continuous spectra.
7. Describe the Bohr model of the hydrogen atom.
8. Define nucleon and isotopes.
9. Explain mass defect in terms of mass-energy equivalent.
10. Explain how the binding energy curve shows that fission and fusion can release energy.
11. Explain the concept of half-life.
12. Explain how alpha and beta decay can stabilize a nucleus.
13. Describe fission reaction and fusion reaction.
14. Explain a chain reaction and nuclear energy.

Survey the LIFEPAAC. Ask yourself some questions about this study. Write your questions here.

The last page of this LIFEPAAC contains formulas and constants that may be used for problem-solving, both on activities and on tests.

I. QUANTUM THEORY

For many centuries, the primary concern of physics was to describe the behavior of large objects. Newtonian mechanics describes the behavior of matter influenced by gravitational fields or by electromagnetic interactions. Newtonian physics had been successful; so successful, in fact, that most physicists believed they were approaching the ultimate description of nature. They thought that in just a matter of time they would be able to work out the details of a perfect, ordered world. Many believed that the future work of physics would be essentially just measuring the last decimal place.

Problems with this neat and supposedly complete theoretical description began to arise about the end of the nineteenth century. More precise experiments opened a whole new world to

the physicist who was beginning to investigate the properties of very small particles. The behavior of these microscopic systems could not be explained in terms of the classical physics and Newtonian mechanics, which had been the physicists' stock and trade up to this point.

In classical physics the physicist had explained the behavior of objects in terms of direct and compelling deductions from a well defined set of experiments. Attempts to do the same from new observations were not successful. The new physics, or *quantum physics*, was too abstract to make deductive explanations possible. The main reasons were twofold: The basic constructs of this quantum theory were removed from everyday experience, and the microscopic particles could not be observed directly.

SECTION OBJECTIVES

Review these objectives. When you have completed this section, you should be able to:

1. Explain when wave theory is applicable and when quantum theory is applicable.
2. Calculate the energy of photoelectrons and X rays.
3. Explain de Broglie waves.
4. State and apply the uncertainty principle.
5. Define emission spectra and absorption spectra.
6. Define line spectra and continuous spectra.
7. Describe the Bohr model of the hydrogen atom.

VOCABULARY

Study these words to enhance your learning success in this section.

absorption spectrum	energy levels	proton
alpha particle	excited atom	quanta
continuous spectrum	ground state	quantum number
de Broglie wave	line spectrum	radiation
electron volt	photoelectric effect	uncertainty principle
emission spectrum	photon	X ray


Note: All vocabulary words in this LIFEPAAC appear in **boldface** print the first time they are used. If you are unsure of the meaning when you are reading study the definitions given.

ELECTROMAGNETIC RADIATION

The wave theory of light was well recognized by the end of the nineteenth century. Experimental work was being conducted in which both light rays and high speed electrons were allowed to strike a metal surface. The results of these experiments

could not be explained by existing theory. As a result a new theory was developed, called the *quantum theory*. At the time, physicists did not fully realize what was happening; but modern physics had been born.

Photoelectric effect. During the late 1800's several experiments disclosed that electrons were emitted when light fell on a metal plate. This phenomenon is the **photoelectric effect**. At first no one was surprised: Light waves were known to carry energy; some of this energy, therefore, could somehow be absorbed by electrons to give them enough kinetic energy to escape from the metal surface.



Soon some problems developed: *First*, the energy of the photoelectrons was not related to the *intensity* of the light. When the light was made brighter, more electrons were emitted; but they all had the same average energy as those emitted by dim light. *Second*, the energy of the photoelectrons was related to the *frequency* of the light. If the frequency was below a certain value, no electrons were emitted even if the light was intense. Conversely, if the frequency was above the value, electrons were emitted even in very dim light.

Photoelectrons emitted by any given frequency were observed to have energy that ranged from zero to some maximum value. If the frequency of the light was increased, the maximum electron energy increased. In other words, a faint blue light produced electrons with higher energy than did a bright red light even though the bright red light produced more electrons. The relationship between KE_{\max} , the maximum photoelectron energy, and the frequency of the incident light can be stated:

$$\begin{aligned} &\text{maximum} \\ &\text{photoelectron} \\ &\text{energy} = h (\text{frequency} - \text{frequency}_0) \\ &KE_{\max} = h (\nu - \nu_0). \end{aligned}$$

The symbol ν (the Greek *nu*) represents the frequency of incident light. The value of h was always the same value, but ν_0 changed for each metal.

An explanation of this phenomenon was given in 1905 by Albert Einstein. He used a result of earlier work by the German physicist Max Planck, who described the radiation emitted by a glowing object. Planck could get his theory to work if he assumed that the light was emitted in little "lumps" of energy. He called these little bursts of energy **quanta** and showed that they had energy of

$$E = h\nu,$$

where ν is the frequency and h is a constant known as *Planck's constant*. The value of Planck's constant h is given as

$$h = 6.63 \cdot 10^{-34} \text{ joule}\cdot\text{sec}.$$

Few physicists really believed in Planck's quanta, but Einstein pointed out that the photoelectric effect was experimental verification of Planck's equation. He suggested that the photoelectric equation be rewritten:

$$h\nu = KE_{\max} + h\nu_0,$$

and be stated:

quantum (incident photon) energy = maximum electron energy + energy required to eject an electron from the metal surface.

A large portion of the photoelectron population will have less than maximum energy for the following reasons:

1. The photon (the quantum of incident light) may have shared its energy with more than one electron.
2. The electron may have lost part of its energy before it got out of the metal.

The energy of photons is so small that the joule is cumbersome. A unit of energy suited to the atomic scale is the **electron volt**. One electron volt is the energy change realized by an electron moving through a potential change of one volt. Numerically,

$$1 \text{ eV} = 1.6 \cdot 10^{-19} \text{ joules}$$

LIFEPACs 1204 and 1205 dealt with diffraction and interference. The observed phenomena could be explained in terms of the wave theory of light. Now you have seen an argument that light behaves like a series of packets of energy, called *photons*, or *quanta*, each **photon** small enough to interact with a single electron. The wave theory cannot explain the photoelectric effect and the quantum theory cannot explain interference and diffraction. Which is the correct theory? Many times before in physics, one theory has been replaced by a new theory; but this apparent conflict was the first time that two different

theories were needed to describe a phenomenon. Developing a model for light is a situation in which science must acknowledge that nature is not an “open book” to man. These two theories are complementary: One is correct in some experiments, but the other theory must be used in others.

Interestingly, light can exhibit either a wave or a particle nature, but never exhibits both in any single situation. Since we cannot reconcile the two descriptive models, we have no choice but to accept them both.



Answer these questions.

1.1 What is the photoelectric effect?

1.2 Why do photoelectrons have a maximum energy?

1.3 Why does faint light not appear as a series of flashes?

1.4 Why can the photoelectric effect not be explained without quantum theory?



Complete these sentences.

1.5 The phenomenon that describes the emission of electrons from a metal surface when light shines on it is the _____ .

1.6 The maximum energy of photoelectrons a. _____ with frequency and b. _____ with increasing intensity of the light. (increases, decreases, remains constant)

1.7 Photon energy is proportional to _____ .

1.8 Photons are also known as _____ .



Solve these problems.

1.9 A yellow lamp emits light with a wavelength of $6 \cdot 10^{-7}$ m.

a. Write the common value for the speed of light, in meters per second. _____